

# Effects of Land Development and Season on Heavy Metal Concentrations in Urban Streams

Emmy Daigle  
August 24, 2018

# Stormwater runoff and pollutants



Hazards of Stormwater Runoff | Hurricane Tips | Hurricane Awareness. (2013, July 17). Retrieved August 23, 2018, from <https://www.national-hurricane-center.org/uncategorized/hazards-of-stormwater-runoff>




# Green infrastructure



Green Streets: Restoring Rivers, Revitalizing Neighborhoods, and Making Streets Safer. (n.d.). Retrieved August 23, 2018, from <https://www.rivernetwork.org/case-study/green-streets-restoring-rivers-revitalizing-neighborhoods-making-streets-safer/>




# Methodology





# Monitoring locations and impervious surfaces




# Results - Season

## Percent difference between seasons - Calcium

Greatest: 25.04% ( $p < 0.001$ )  
(Moderate development, GI present)

Lowest: 15.99% ( $p = 0.007$ )  
(Moderate development, GI absent)




Moderate development, GI present

# Results - Season

Percent difference between  
seasons - Magnesium

Greatest: 28.8% ( $p = 0.040$ )  
(Low development, GI present)

Lowest: 19.5% ( $p = 0.001$ )  
(Moderate development, GI absent)




Low development, GI present

# Results - Season

Percent difference between  
seasons - Hardness

Greatest: 25.7% ( $p < 0.001$ )  
(Moderate development, GI present)

Lowest: 17.3% ( $p = 0.003$ )  
(Moderate development, GI absent)



Moderate development, GI present




# Results - Development

Percent difference between  
low and high development  
levels - Calcium

Greatest: 34.17 ( $p < 0.001$ )  
(Wet season, GI present)

Lowest: 27.31% ( $p = 0.010$ )  
(Dry season, GI present)




Wet season, GI present

# Results - Development

Percent difference between  
low and high development  
levels - Magnesium

Greatest: 35.11% ( $p < 0.001$ )  
(Dry season, GI absent)

Lowest: 24.75% ( $p = 0.021$ )  
(Dry season, GI present)




Dry season, GI absent

# Results - Development

Percent difference between  
low and high development  
levels - Hardness

Greatest: 33.54% ( $p < 0.001$ )  
(Wet season, GI present)

Lowest: 26.25% ( $p = 0.013$ )  
(Dry season, GI present)





Wet season, GI present





# Results - Green infrastructure



# Results - Summary: Calcium





# Results - Summary: Magnesium





# Results - Summary: Hardness



# Conclusions and Future areas of study



Water and Wellness: Green Infrastructure for Health Co-Benefits. (n.d.). Retrieved August 24, 2018, from <http://stormwater.wef.org/2014/04/water-wellness/>

# Acknowledgements

The REU Site is supported by the National Science Foundation under grant no 1758006.

This research was also supported by NSF #1444755. Views expressed are our own and do not necessarily reflect the views of NSF.

This research was directed and overseen by PSU Geography Department Chair, Heejun Chang.

Precipitation data was provided by the United States Geological Survey.

Water quality data was collected and provided by City of Portland BES.

GIS analysis was provided by Josh Spector and Emma Brenneman.

